


CERTIFICATE OF MAILING (37 CFR 1.8a)

I hereby certify that this paper (along with any paper referred to as being transmitted therewith) is being deposited with the United States Postal Service on the date shown below with sufficient postage as first class mail in an envelope addressed to the: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

Dennis P. Tramaloni

Date: September 17, 2007


(Signature)

PATENT APPLICATION

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:

Group No.: 1751

Pierre Bartsch, et al.

Serial No.: 10/594,162

Filed: September 26, 2006

For: PHARMACEUTICAL COMPOSITIONS OF PYRIMIDINE-2,4,6-TRIONES

TRANSMITTAL OF CERTIFIED COPY

September 17, 2007


Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Dear Sir:

Attached please find the certified copy of the foreign application from which priority is claimed for this case:

<u>Country</u>	<u>Application No.</u>	<u>Filing Date</u>
Europe	04007921.2	April 1, 2004

Respectfully submitted,


Dennis P. Tramaloni
Attorney for Applicant
Reg. No. 28,542
Hoffmann-La Roche Inc.
340 Kingsland Street
Nutley, New Jersey 07110
Phone: (973) 235-4475

DPT/ks
Enclosure
316446



**Europäisches
Patentamt**

**European
Patent Office**

**Office européen
des brevets**

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

04007921.2

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk



Anmeldung Nr:
Application no.: 04007921.2
Demande no:

Anmeldetag:
Date of filing: 01.04.04
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

F. HOFFMANN-LA ROCHE AG

4070 Basel
SUISSE
Université de Liège
25, quai Van Beneden
4020 Liège
BELGIQUE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Pharmaceutical compositions of pyrimidine-2,4,6-triones

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

A61K47/48

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PL PT RO SE SI SK TR LI

Pharmaceutical compositions of pyrimidine-2,4,6-triones

The invention comprises a pharmaceutical composition of pyrimidine-2,4,6-triones (trioxypyrimidines), methods for the manufacture and uses thereof.

Matrix metalloproteases (MMPs) are a family of zinc- and calcium-dependent proteases that are capable of degrading the extracellular matrix (ECM) and basement membrane (Egeblad, M., and Werb, Z., Nat. Rev. Cancer 2 (2002) 161-174; Overall, C.M., and Lopez-Otin, C., Nat. Rev. Cancer 2 (2002) 657-672). They are believed to have pivotal roles in embryonic development and growth (Holmbeck, K., et al., Cell 99 (1999) 81-92; Vu, T.H., et al., Cell 93 (1998) 411-422) as well as in tissue remodeling and repair (Shapiro, S.D., Curr. Opin. Cell Biol. 10 (1998) 602-608; Lund, L.R., et al., EMBO J. 18 (1999) 4645-4656). Excessive or inappropriate expression of MMPs may therefore contribute to the pathogenesis of many tissue-remodelling processes, including tumor progression (Egeblad, M., and Werb, Z., Nat. Rev. Cancer 2 (2002) 161-174; Overall, C.M., and Lopez-Otin, C., Nat. Rev. Cancer 2 (2002) 657-672) and aneurysm formation (Carmeliet, P., et al., Nat. Genet. 17 (1997) 439-444). MMP effects are far from being restricted to ECM degradation (Chang, C., and Werb, D., Trends Cell Biol. 11 (2001) S37-43). Peptide growth factors that are sequestered by ECM proteins become available once degraded by MMP-9 (Manes, S., et al., J. Biol. Chem. 274 (1999) 6935-6945). MMPs can increase the bioavailability of VEGF (Bergers, G., et al., Nat. Cell Biol. 2 (2000) 737-744) but also generate angiogenesis inhibitors such as angiostatin by cleavage of plasminogen (Dong, Z., et al., Cell 88 (1997) 801-810). MMPs are thought to be involved in the mobilization of bone marrow stem cells (Janowska-Wieczorek, A., et al., Blood 93 (1999) 3379-3390). High concentration of MMP9 was observed during the G-CSF induced HPC mobilization (Carstanjen, D., et al., Transfusion 42 (2002) 588-596).

Trioxypyrimidines are compounds from a well-known structural class. Such compounds are described in, for example, US Patent Nos. 6,242,455 and 6,110,924; WO 97/23465; WO 98/58915; WO 01/25217, which are incorporated herein by reference, and Grams, F., et al., Biol. Chem. 382 (2001) 1277-1285, and are effective and highly selective for MMP-2, MMP-9 and MMP-14.

Cyclodextrins are cyclic carbohydrates derived from starch. They differ from one another by the number of glucopyranose units in their structure. The parent cyclodextrins contain six, seven and eight glucopyranose units, and are referred to as alpha, beta and gamma cyclodextrins respectively. The α -, β - or γ -cyclodextrins prepared by enzymatic starch conversion differ in the diameter of their hydrophobic cavity and are generally suitable for the inclusion of numerous lipophilic substances.

Trioxypyrimidines which are highly potent MMP inhibitors are only poorly soluble in water and water-based solvents. The object of the invention is therefore to provide an aqueous composition in which such a trioxypyrimidine is soluble and whereas such an aqueous composition of such a trioxypyrimidine can be used as a pharmaceutical composition.

Summary of the Invention

It was surprisingly found that a trioxypyrimidine-cyclodextrin complex formed of a trioxypyrimidine derivative represented by the below-described formula (I) and a water-soluble cyclodextrin (further abbreviated as CD) exhibits enhanced water solubility, excellent stability, and low topical stimulation and is useful as a therapeutic agent.

Accordingly, the present invention provides a trioxypyrimidine-cyclodextrin complex formed of a trioxypyrimidine derivative or a salt thereof and a cyclodextrin, preferably α -, β - or γ -cyclodextrin or a water-soluble cyclodextrin derivative (water-soluble being defined as a solubility of at least 0.5 gr/100ml water at 25°C), wherein the trioxypyrimidine derivative is represented by formula (I).

Such a complex according to the invention is an inclusion complex of trioxypyrimidine-cyclodextrin and is provided in a liquid or solid form.

In the complex according to the present invention, preferably 1 mol of trioxypyrimidine is complexed and enclosed by about 1 mol to 2 mol of cyclodextrin, preferably of β - or γ -cyclodextrin or a derivative thereof.

The present invention also provides a pharmaceutical agent for the treatment of a patient in the need thereof, preferably for the treatment of bronchial inflammatory

diseases, containing a trioxypyrimidine -cyclodextrin complex according to the invention as an active component in a pharmaceutical effective amount.

5 The pharmaceutical agent according to the invention is applicable therapeutically, prophylactically or preventively, to pathologies resulting from a very important or unsuitable MMP expression. Preferably such treatment is a therapeutic, prophylactic or preventive treatment of rheumatoid arthritis, tumors, metastatic invasion, osteoporosis, macular degeneration, diabetic retinopathies, ulcerations of the cornea, atherosclerosis, bronchial inflammatory diseases, bronchial inflammatory diseases such as asthma, chronic obstructive pulmonary disease or
10 emphysema.

The present invention also provides an injection formulation containing a trioxypyrimidine-cyclodextrin complex according to the invention in a pharmaceutically effective amount.

15 A further object of this invention is a liquid aqueous formulation of a complex according to the invention, the pharmaceutically acceptable carrier is water, the composition to administrate being an aqueous solution. The active substance according to the invention is then in the complex state by inclusion in a cyclodextrin in solution in water.

20 A further object of this invention is a complex according to the invention in a solid state, the complex is in the form of a powder dissolvable in water and to dissolve before administration or to administrate on its own.

25 A further object of this invention is a complex included in different galenical forms according to the desired form of administration which can be tablets, capsules, multiparticulate systems, oral solutions, oral suspensions, solutions, suspensions, and implants for parenteral administration, solutions or powders for inhaling, hydrophilic or lipophilic type creams and ointments, aqueous or hydro-alcoholic gels, lotions, for topical, transcutaneous or vaginal use, intra-uterine devices, solutions, suspensions, implants, for ophthalmic use, suppositories, suspensions, sprays, solutions, and foams for rectal use.

30 The present invention further provides use of such a pharmaceutical agent in a pharmaceutically effective amount for the treatment of such diseases in a patient

suffering from such a disease, preferably bronchial inflammatory diseases. The complex according to the invention is preferably administrated at a topical, percutaneous, transdermal, oral or parenteral level.

5 The present invention further provides a method for the manufacture of a pharmaceutical agent, preferably for the treatment of such diseases, preferably bronchial inflammatory diseases, characterized by complexing a trioxypyrimidine with cyclodextrin in a pharmaceutically effective amount in water or buffered aqueous solution preferably containing, in addition, an auxiliary substance, buffer, preservative, solvent and/or viscosity modulating agent.


10 The preferred cyclodextrins are

- alpha-cyclodextrin and its synthetic derivatives such as HP α CD, methylated α CD, hydroxybutyl α CD, maltosyl α CD, glucosyl α CD.
- beta-cyclodextrin and its synthetic derivatives such as HP β CD, SBE β CD, RM β CD, DIME β CD, TRIME β CD, hydroxybutyl β CD, glucosyl β CD, maltosyl β CD.
- 15 - gamma-cyclodextrin and its synthetic derivatives such as HP γ CD, RM γ CD and DIME γ CD, hydroxybutyl γ CD, glucosyl γ CD, maltosyl γ CD.

20 This invention also concerns use of a pharmaceutical composition including, in a therapeutically effective quantity, a pyrimidine-2,4,6-trione and at least one cyclodextrin, as well as possibly a pharmaceutically acceptable carrier, for the manufacture of a medicine for a therapeutic, prophylactic or preventive treatment of the above-mentioned illnesses.

Detailed Description of the Invention

Pyrimidine-2,4,6-triones (trioxypyrimidines) according to the present invention are those of formula (I)



wherein

5 R^1 is C_3 - C_{20} alkyl, which may optionally be interrupted once or several times by -S-, -O- or -NH-; or

a group W-V, wherein

W is a chemical bond or phenyl; and

10 V is phenyl, phenyloxy, phenylthio, phenylsulfinyl, phenylsulfonyl or phenylamino, which phenyl moieties may be unsubstituted or substituted once or several times by halogen, hydroxy, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_1 - C_6 -alkylthio, C_1 - C_6 alkylsulfinyl, C_1 - C_6 -alkylamino, cyano, nitro or C_1 - C_6 -alkylsulfonyl; and

15 R^2 is C_1 - C_{10} alkyl, which alkyl group is unsubstituted or substituted one or two times by hydroxy or amino and may optionally be interrupted once or several times by -S-, -O- or -NH-;

20 a benzoyl group, which may be unsubstituted or substituted once or several times by halogen, hydroxy, nitro, C_1 - C_6 -alkoxy, C_1 - C_6 -alkylamino, C_1 - C_6 -alkylthio, C_1 - C_6 -alkylsulfinyl, C_1 - C_6 -alkylsulfonyl,

amidosulfonyl, C₁-C₆-alkylamidosulfonyl, bis-C₁-C₆-alkylamido-sulfonyl;

a heteroaromatic acyl group; or

5 a phenyl- or heteroaryl group, which are unsubstituted or substituted once or several times by halogen, hydroxy, C₁-C₆-alkoxy, C₁-C₆-alkylamino, C₁-C₆-dialkylamino, cyano, C₁-C₆-alkyl, C₂-C₆ alkenyl, C₂-C₆-alkinyl, C₁-C₆-acyl, C₁-C₆-alkylthio, C₁-C₆-alkylsulfonyl, C₁-C₆-alkylsulfinyl, C₁-C₆-alkylaminocarbonyl, aminocarbonyl, C₁-C₆-alkylamidosulfonyl, amidosulfonyl, bis-C₁-
10 C₆-alkylamidosulfonyl, nitro, C₁-C₆-alkoxycarbonyl, carboxy.

An object of the present invention is the use of the compounds of formula (I), as well as their pharmaceutically acceptable salts, enantiomeric forms, diastereoisomers and racemates, in the manufacture of novel pharmaceutical preparations.

15 As used herein for R¹, the term „C₃-C₂₀ alkyl“ represents a linear or a branched saturated hydrocarbon containing from 3 to 20-, preferably from 4 to 12- and more preferably from 8 to 12 carbon atoms. Examples are butyl, hexyl, octyl, decyl, 2-ethylhexyl, 2 ethyloctyl. Preferred C₃-C₂₀ alkyl residues are n-octyl and n-decyl. The C₃-C₂₀ alkyl group may be interrupted once or several times by -S-, -O- or -
20 NH-, preferably by -O-. Examples for such C₃-C₂₀ alkyl groups are 5-ethoxy-n-pentyl, 9-methoxy-n-octyl.

The substituents in the phenyl moieties of „V“ are preferably located in p- and/or meta- position.

25 Preferably the group „W-V“ is p-butoxyphenyl, biphenyl, phenoxyphenyl, p-chloro-phenoxyphenyl, p-bromo-phenoxyphenyl, 3,4 dichloro-phenoxyphenyl.

The term „C₁-C₁₀-alkyl“ as used in R² represents a linear or branched saturated hydrocarbon, containing from 1 to 10, preferably from 1 to 6 and more preferably from 1 to 4 carbon atoms. Said C₁-C₁₀-alkyl may be interrupted once or several times by -S-, -O- or -NH-, preferably by -O- and more preferably in such a way to
30 give a group which is composed of ethyleneoxy fragments. Preferred examples of

C₁-C₁₀-alkyl groups are hydroxyethyl; hydroxypropyl; ethoxyethyl; 1,2-bisethoxyethyl; 1,2-bis-hydroxy-ethyl.

5 The term heteroaromatic as used in „heteroaromatic acyl group“ in R² denotes a five- or six membered aromatic ring, wherein one, two or three ring atoms are oxygen, nitrogen or sulfur, and the remaining ring atoms being carbon atoms. Said heteroaromatic group may be fused to another phenyl ring. Examples for such heteroaromatic acyl groups are furanecarboxyl, thiophenecarboxyl, 4-imidazolylcarboxyl, 3-benzthiophenecarboxyl, pyridylcarboxyl. Preferred examples are furanecarboxyl and thiophenecarboxyl.

10 The term „heteroaryl“ as used herein means heteroaromatic as defined above. Preferred heteroaryl groups are electron deficient residues such as the nitrogen containing 6-membered rings like pyridine, pyrimidine, pyrazine or 1,3,5-triazine. Especially preferred are the heteroaryl groups pyrimidinyl or pyrazinyl.

15 Substituents which may be present on the phenyl or heteroaryl groups of R² are principally located at any position suitable for the respective substitution reaction. Preferably one or two substituents are present in para and/or meta position.

20 The term „C₁-C₆-alkyl“ as used herein alone or in combination with C₁-C₆-alkoxy, C₁-C₆-alkylamino, C₁-C₆-dialkylamino, C₁-C₆-acyl, C₁-C₆-alkylthio, C₁-C₆-alkylsulfonyl, C₁-C₆-alkylsulfinyl, C₁-C₆-alkylaminocarbonyl, C₁-C₆-alkylamidossulfonyl, bis-C₁-C₆-alkylamidossulfonyl or C₁-C₆-alkoxycarbonyl denotes a linear or branched, saturated hydrocarbon with 1 to 6-, preferably 1 to 4 carbon atoms. Preferred examples are methyl, ethyl, propyl, isopropyl or tert.-butyl.

25 The term „C₂-C₆-alkenyl“ as used herein denotes a linear or branched unsaturated hydrocarbon containing 2 to 6-, preferably 2 to 5 carbon atoms and one or two double bonds. If two double bonds are present they can be isolated- or conjugated double bonds, preferably conjugated double bonds. Preferred examples are allyl or pentadienyl.

30 The term „C₂-C₆-alkinyl“ as used herein denotes a linear or branched hydrocarbon containing 2 to 6-, preferably 2 to 4 carbon atoms. The preferred example is propargyl.

The term „halogen“ means fluorine, chlorine, bromine, iodine, preferably chlorine or bromine.

The term „several times“ as used herein means one, two, three or four times, preferably one or two times.

5 The term “pharmaceutically acceptable salt” as used herein before refers to
conventional acid-addition salts or base-addition salts that retain the biological
effectiveness and properties of the compounds of formula (I) and are formed from
suitable non-toxic organic or inorganic acids or organic or inorganic bases. Sample
10 acid-addition salts include those derived from inorganic acids such as hydrochloric
acid, hydrobromic acid, hydroiodic acid, sulfuric acid, sulfamic acid, phosphoric
acid and nitric acid, and those derived from organic acids such as p-toluenesulfonic
acid, salicylic acid, methanesulfonic acid, oxalic acid, succinic acid, citric acid, malic
15 acid, lactic acid, fumaric acid, and the like. Sample base-addition salts include
those derived from ammonium, potassium, sodium and, quaternary ammonium
hydroxides, such as for example, tetramethylammonium hydroxide. The chemical
modification of a pharmaceutical compound (i.e., a drug) into a salt is a technique
well known to pharmaceutical chemists to obtain improved physical and chemical
20 stability, hygroscopicity, flowability and solubility of compounds (see, e.g., Ansel,
H., et. al., Pharmaceutical Dosage Forms and Drug Delivery Systems, 6th ed.,
(1995), pp. 196 and 1456-1457).

The compounds of the present invention can be prepared as described in
EP 0 869 947 and WO 01/25217.

According to the invention, the following compounds are particularly preferred:

25 5-Biphenyl-4-yl-5-[4-(4-nitro-phenyl)-piperazin-1-yl]pyrimidine-2,4,6-trione
(Compound I)

5-(4-Phenoxy-phenyl)-5-(4-pyrimidin-2-yl-piperazin-1-yl)-pyrimidine-2,4,6-
trione
(Compound II)

5-[4-(4-Chloro-phenoxy)-phenyl]-5-(4-pyrimidin-2-yl-piperazin-1-yl)-
pyrimidine-2,4,6-trione
(Compound III)

5 5-[4-(3,4-Dichloro-phenoxy)-phenyl]-5-(4-pyrimidin-2-yl-piperazin-1-yl)-
pyrimidine-2,4,6-trione
(Compound IV)

5-[4-(4-Bromo-phenoxy)-phenyl]-5-(4-pyrimidin-2-yl-piperazin-1-yl)-
pyrimidine-2,4,6-trione
(Compound V).

10 It is also apparent that when the trioxypyrimidine derivative (I) contains an acidic moiety such as a carboxylic group or a sulfonyl group, the derivative can form a salt with a base via the acidic moiety.

15 In addition to the above-described adduct-type salt, the trioxypyrimidine may take a hydrate form or a solvated form. The hydrate and the solvate include both that of the free compound of the formula (I) and a salt of the compound of the formula (I). They also include a tautomer of the compound of the formula (I).

20 Cyclodextrins (CD) according to the invention are cyclic oligosaccharides produced by enzymatic degradation of starch, which are composed of a variable number of glucopyranose units, mostly 6, 7 or 8: these cyclodextrins are respectively named α , β , and γ cyclodextrins (α CD, β CD and γ CD). Cyclodextrins according to the invention are cyclodextrins per se or cyclodextrin derivatives, which are at least water soluble in an amount of 0.5gr/100ml at 25°C.

25 The water-soluble cyclodextrin preferably used in the present invention refers to a cyclodextrin having water solubility of at least that of β -cyclodextrin. Examples of such water-soluble cyclodextrin are sulfobutylcyclodextrin, hydroxypropyl-cyclodextrin, maltosylcyclodextrin, and salts thereof. In particular, sulfobutyl- β -cyclodextrin, hydroxypropyl- β -cyclodextrin, maltosyl- β -cyclodextrin, and salts thereof.

30 Cyclodextrins preferred according to the invention are also methylcyclodextrins (products of the cyclodextrins methylation), dimethylcyclodextrins (DIMEB)

(preferably substituted in 2 and in 6), trimethylcyclodextrins (preferably substituted in 2, 3 and 6), "random methylated" cyclodextrins (preferably substituted at random in 2, 3 and 6, but with a number of 1,7 to 1,9 methyl by unit glucopyranose, RM β CD), hydroxypropylcyclodextrins (HPCD, 5 hydroxypropylated cyclodextrins preferably substituted randomly mainly in position 2 and 3 (HP- β CD, HP- γ CD)), sulfobutylethercyclodextrins (SBECD), hydroxyethyl-cyclodextrins, carboxymethylethylcyclodextrins, ethylcyclodextrins, amphiphilic cyclodextrins obtained by grafting hydrocarbonated chains in the hydroxyl groups and being able to form nanoparticles, cholesterol cyclodextrins 10 and triglycerides-cyclodextrins obtained by grafting cyclodextrins monoaminated (with a spacer arm).

The trioxypyrimidine -cyclodextrin complex of the present invention may be obtained by producing an aqueous solution containing the trioxypyrimidine or a salt thereof and a water-soluble cyclodextrin. The water-soluble cyclodextrin is used 15 in an amount of preferably one mol or more based on 1 mol per mol trioxypyrimidine or a salt thereof, more preferably 1-10 mol, and particularly preferably 1-2 mol cyclodextrin per mol trioxypyrimidine.

The higher the concentration of the water-soluble cyclodextrin, the more the solubility of the trioxypyrimidine increases. No particular limitation is imposed on 20 the method for producing the aqueous solution, and for example it is produced by use of water or a buffer in a temperature range approximately from -5 to 35° C.

When a cyclodextrin aqueous solution is stirred with an excess of a trioxypyrimidine of formula I, there is a complex formation between these two molecules. Reaching the equilibrium takes, however, about at least a few days, so 25 that after a few hours or even after one day, the improved solubility of trioxypyrimidines according to the invention is not found. The filtration of the solution allows recovering the complex in solution in the filtrate, the complex being soluble in water. The complex can also be obtained by mixing a solubilized known quantity of a trioxypyrimidine of formula I in aqueous solution with a solubilized 30 known quantity of CD by calculating the adequate proportions.

Another way of obtaining a complex is to add a solution of a trioxypyrimidine of formula I in a solvent (e. g. alcohol, acetone, etc) to a cyclodextrin aqueous

solution. The complex can be formed after sufficient stirring, either after evaporation of the solvent, or even in the presence of the solvent.

5 The lyophilization or the nebulization of solutions of the complex according to the invention allows the complex to be obtained in solid form. One can thus obtain a complex in the form of an amorphous powder. It is also possible to obtain the complex in the solid state after dissolution of CD and a trioxypyrimidine of formula I in an appropriate organic solvent and further evaporation of the solvent.

10 Other methods can be used for solid complexes preparation which are violent stirring of a suspension of a trioxypyrimidine of formula I and CD in a very small quantity of water, then complex collecting after drying or the use of CO₂ in a supercritical state for mixing a trioxypyrimidine of formula I and CD in presence of CO₂ in a supercritical state.

15 The complex according to the present invention can be prepared, for example, in a manner known per se from a solution or using the paste method, where the weight ratio of cyclodextrin to trioxypyrimidine should be between 2 (2:1) to 540 (540:1), and is preferably between 2 to 25, particularly preferably in the region of 2.6 to 3.5 (for a 1:1 complex with cyclodextrin) or of 5.2 to 6.2 (for a 1:2 complex with cyclodextrin) for a molecular weight of the cyclodextrin of about 1,300.

20 It is preferred to prepare the complex from a concentrated, aqueous cyclodextrin preparation. The cyclodextrin concentration of the preparation is preferably between 50 and 400 mM. Preference is given to a cyclodextrin concentration of from 100 to 250 mM. Depending on the consistency, the mixtures are intensively stirred or kneaded. The percent by weight of the cyclodextrin is based upon the total weight of the aqueous cyclodextrin preparation.

25 The reaction temperature is usually between 20° C and 80° C, preferably between 20° C and 60° C, particularly preferably between 25° C and 45° C. The reaction time depends on the temperature and is at least some days. Preference is given to a reaction time of at least 7 days to reach equilibrium of complex formation. Subsequently, the reaction mixture is filtrated, if undissolved material is still present, or used directly, if completely dissolved. If desired, the complex can be
30 isolated, e.g., by chromatographic means. Preferably, the concentrations and ratio of trioxypyrimidine and cyclodextrin are such that complex formation has occurred

completely (reached the equilibrium) and no undissolved or uncomplexed trioxypyrimidine is detectable.

5 According to the invention it has been established that complexes between a trioxypyrimidine of formula I and a cyclodextrin increase the solubility of the trioxypyrimidine in water amazingly. It was also found that the formation of the complex did not interfere with the pharmacological properties of the trioxypyrimidine.

10 All these properties allow to prepare liquid formulations as solutions for injection or for nebulization and allow the bioavailability to be improved, in particular orally. A trioxypyrimidine-cyclodextrin complex of the present invention may be used as such or in a powder form which is obtained by removing co-existing water. Examples of the method for removing water include lyophilization and drying under reduced pressure. A powder product obtained from lyophilization is particularly preferred.

15 The trioxypyrimidine -cyclodextrin complex of the present invention exhibits its effects through either oral administration or parenteral administration, and it is preferably formed into a formulation for parenteral administration, particularly an injection formulation or topical administration, particularly an aerosol formulation.

20 The dose of the complex of the present invention may be modified appropriately in accordance with the age, body weight, and severity of the patient's symptom and the complex may be administered at a single time or in a divided manner. Examples of the form of formulation include tablets, capsules, powders, and granules. These may be produced through a known technique by use of typical additives such as
25 excipients, lubricants, and binders.

The invention relates to a method used for treating bronchial inflammatory diseases in a host mammal in need of such treatment, e.g., especially asthma and chronic obstructive pulmonary disease (COPD) by the application of a complex according to the invention to a patient in a pharmaceutically effective amount.
30 Asthma is an inflammatory disease of the bronchial tree related or not to an allergen exposure. This inflammation provokes symptoms in patients by stimulating the bronchial smooth muscles to contract, enhancing the mucus

secretion, and inducing bronchial morphological changes thought to be an aggravating factor regarding the course of the disease. Airway hyperresponsiveness is a hallmark of the disease and is responsible for most of symptoms. Bronchial tree is a very complex tissue with many cell types (epithelial cells, smooth muscle cells, inflammatory cells, nerves, mucus producing cells, fibroblasts, and the like) and the bronchial remodelling events which comprise many aspects mainly consist in a deposition of extracellular matrix components in the bronchial walls and an hyperplasia of the mucus producing cells. The use of complexes according to the invention inhibits the inflammatory cells influx in the compartments of bronchoalveolar lavage and peribronchial tissue and inhibits the hyperresponsiveness defined as an abnormal response to stimulating agents such as methacholine. The disease and current treatments are reviewed in e.g.: GINA Workshop Report, Global Strategy for Asthma Management and Prevention (NIH Publication No. 02-3659).

The invention therefore further relates to a method for treating or preventing in a host mammal in need of such treatment chronic obstructive pulmonary diseases using complexes according to the invention. In such a disease, bronchi are inflamed and the mucus glands are hyperplastic and produce high amounts of mucus. The bronchial wall is abnormal and deposition of abnormal extracellular matrix components increases the resistance to airflow. The disease and current treatments are described by, e.g., Fabbri, L.M., and Hurd, S.S., Eur. Respir. J. 22 (2003) 1–2.

The invention therefore further relates to a method for treating or preventing in a host mammal in need of such treatment emphysema using complexes according to the invention. In such a disease, the alveolar walls are destroyed by proteolytic processes and this destruction impairs the transfer of oxygen to the blood. Physiological problems also occurs because of the derived hyperinflation which causes abnormalities in the ventilation by causing a dysfunction of respiratory muscles and because of a hypertension in pulmonary arteries leading to cardiac failure in advanced stages.

According to the invention the trioxypyrimidine-cyclodextrin complexes are preferably administered over several months or years, to the patient in need of such a therapy. The complexes are administered preferably by the aerosolization of a liquid or powder formulation, with non toxic doses ranging between micro and nanomolar concentrations per kg and day.

The exact dosage of the complexes according to the invention will vary, but can be easily determined. In general, the daily dosage of the complexes will range between 1 μ mol/kg and day to 100 nmol/kg and day (concentration of the trioxypyrimidine in the complex).

5 The pharmaceutical compositions are preferably aqueous compositions having physiological compatibility. The compositions include preferably, in addition, a pharmaceutically acceptable additive such as buffer, preservative and/or auxiliary substance. Appropriate buffer systems are based on sodium phosphate, sodium acetate or sodium borate. Preservatives are required to prevent microbial
10 contamination of the pharmaceutical composition during use. Suitable preservatives are, for example, benzalkonium chloride, chlorobutanol, methylparabene, propylparabene, phenylethyl alcohol, sorbic acid. Such preservatives are used typically in an amount of 0.01 to 1% weight/volume.

15 Suitable auxiliary substances and pharmaceutical formulations are described in Remington's Pharmaceutical Sciences, 16th ed., 1980, Mack Publishing Co., edited by Oslo et al. Typically, an appropriate amount of a pharmaceutically-acceptable salt is used in the formulation to render the formulation isotonic. Examples of a pharmaceutically acceptable substances include saline, Ringer's solution and dextrose solution. The pH of the solution is preferably from about 5 to about 8, and
20 more preferably from about 7 to about 7.5.

A preferred formulation according to the invention is an injectable or nebulizable formulation, preferably prepared from CD and trioxypyrimidine in a molar ratio of 1 to 500.

25 The complex is prepared by dissolving CD in water, adding a trioxypyrimidine of formula I and heat in a water bath until the latter is completely dissolved. Preferably the solution is sterilized by filtration. Preferably the solution has a osmolality of 200-400, preferably about 300 mOs/kg. The pH is about 7.2. The concentration of trioxypyrimidine and/or of CD can be modified in function of the requirements. It is preferred to adjust the tonicity by addition of NaCl.

30 A preferred formulation for nebulization contains trioxypyrimidine, CD, NaCl and water. Especially preferred is a combination of (for 200 ml of solution):

Trioxypyrimidine 0.05-0.2 g, preferably 0.1 g; 10-50 g CD, preferably 20 g CD, preferably HP β CD; sodium chloride 1.2-1.5 g, preferably 1.42 g (isotonicity) and water, preferably pyrogen-free, sterile, purified water ad 200 ml.

5 The solution was prepared by dissolving CD in 100 ml of purified water, adding trioxypyrimidine and NaCl by stirring so as to dissolve them and complete with water so as to obtain 200 ml of solution. Preferably the solution is sterilized by filtration through a 0.22 μ m polypropylene membrane or by a steam sterilization process.

10 Other preferred formulations are ophthalmic use formulations, oral use formulations, intra-uterine devices. Associations with other systems can also be considered, like nano- or microparticles or liposomes for example.

15 The following examples, references, and figures are provided to aid the understanding of the present invention, the true scope of which is set forth in the appended claims. It is understood that modifications can be made in the procedures set forth without departing from the spirit of the invention.

Description of the Figures

20 **Figure 1** shows the Compound I solubility obtained for both RM β CD and HP- β -CD. Phase solubility diagrams are both of A_p type which means that CDs form complexes of stoichiometry 1:1 and 1:2. Stability constants were then calculated and their values are given in table 6.

Figure 2 NMR spectrum of the complex of Compound I and DIME β CD (upper part) and of DIME β CD alone (lower part).

25 **Figure 3** NMR spectra of compound I (on the top), of DIME β CD (on the right-hand side) and T-ROESY (in the middle).

Figure 4 Effects of intraperitoneal injection of a Compound I suspension on BAL eosinophil counts (Fig 2a) and peribronchial inflammation score (Fig 2b). Controls are mice exposed only to

PBS and not allergen (PBS) and mice exposed to ova by inhalation and placebo by intraperitoneal injection (OVA).

Figure 5 Therapeutic effects of Compound I-HP- β -CD complex, fluticasone and placebo (PLAC) administered by aerosols on BAL eosinophilia (5a), peribronchial inflammation score (5b), and tissue eosinophils infiltration score (5c) in a short term (5 days) allergen exposure model.

Figure 6 Therapeutic effects of Compound I-HP- β -CD complex, fluticasone and placebo (PBS) administered by aerosols on BAL eosinophilia (6a), peribronchial inflammation score (6b), and tissue eosinophils infiltration score (6c) in a long term (11 weeks) allergen exposure model. Mice sensitized but unexposed to allergens (PBS) and mice sensitized and exposed to OVA (PLAC) were treated by PBS inhalation.

Abbreviations

CD	cyclodextrin
β CD	β -cyclodextrin
γ CD	γ -cyclodextrin
DIME β CD	dimethyl β -cyclodextrin
HP β CD	Hydroxypropyl β -cyclodextrin
RM β CD	random methylated β -cyclodextrin

Example 1

Preparation of a soluble complex of compound I and cyclodextrin (CD)

1.1 Weigh 20 mg of compound I. Add 2 ml of solution of HP β CD 200 mM. Stir for 24 h at 37°C. Filter in Millipore filter Millex HV 0.45 μ m. The solution obtained after filtration contains the complex compound I-CD in solution.

1.2. Weigh 2.5 mg of compound I. Add 2 ml of solution of HP β CD 200 mM. Stir at 37°C for 24 h or until compound I is completely dissolved. The solution obtained in this way contains the complex compound I-CD.

Example 2

Phase solubility studies

At the time of a complex formation, compound I, practically insoluble in water (<0.6 µg/ml, MW: 485), solubilizes dramatically. The increase in the solubility of compound I is thus proof of a complex formation between Compound I and the CD. Complex formation and achievement of the equilibrium is found at 20% after 1 day, 40% after 4 days, and 100% after 7 days. The solubility diagrams (Fig. 1) are carried out by adding an excess of compound I to the CD solutions of increasing concentration. After 7 days of stirring in thermostatically controlled baths at 37°C, these solutions are filtered and the solubilized compound I quantity is dosed by HPLC. The βCD and the γCD as well as their synthetic derivatives have allowed complexes with compound I to be formed. The β-CD and the HPβCD come from ROQUETTE (France), the RMβCD and the γCD have been supplied by Wacker (Germany).

Preparation of CD aqueous solutions:

- βCD: Solutions containing 2, 4, 8, 10, 12, 16 mM.
- HPβCD: Solutions containing 10, 25, 50, 75, 100, 150, 200 mM.
- RMβCD: Solutions containing 10, 25, 50, 75, 100, 150, 200 mM.
- γCD: Solutions containing 10, 25, 50, 100, 75, 150 mM.

Complex formation:

The flasks containing compound I and the cyclodextrins are placed being stirred in thermostatically controlled baths at 37°C for 7 days, so that the complexation balance is reached. After this time, the suspensions are filtered with the help of a milli-pore filter 0.22 µm in PVDF and the filtrate is dissolved in DMSO in the mobile phase to obtain concentration samples situating themselves on the calibrating line. They are then dosed thanks to the validated HPLC method, described below.

Dosage of Compound I: HPLC Method

Equipment:

5 Pump Merck-Hitachi model L-7100, sampler Merck-Hitachi L-7200, furnace Merck-Hitachi L-7350, detector Merck-Hitachi Diode array detector L-7455, interface D-7000, the set being piloted by the data acquirement software "Chromatography Data Station Software" supplied by Merck-Hitachi.

Stationary phase:

Lichrocart Column (125x4 mm d.i.) filled with a stationary phase of octylsilane C8 LiChorspher® 60RP-Select B (5µm) Merck.

10 Chromatographic conditions:

Mobile phase: mixture of phosphate buffer 0,05 M to pH=3 and methanol (30/70, v/v). Gas is extracted by an ultrasound passage for 15 minutes, output: 1 ml/min, λ of U.V. detection: 265 nm, Working temperature: 30°C, Injection volume: 20µl.

15 The results of the HPLC dosages of Compound I are reproduced in the tables below (Tables 1 to 4) for each cyclodextrin. Without cyclodextrin, the solubility obtained is 0,56 µg/ml.

Table 1:

Solubility of Compound I in the presence of HPβCD

Concentration of CD in mM	Concentration of Compound I in µg/ml
2	2.4
4	6.8
8	6.6
10	3.4
12	1.6
16	1.5

Table 2:
Solubility of Compound I in the presence of HP β CD

Concentration of CD in mM	Concentration of Compound I in $\mu\text{g/ml}$
10	187.8
25	213.4
50	235.8
75	1129.3
100	1664.6
150	3106.5
200	4962.1

Table 3:
Solubility of Compound I in the presence of RM β CD

Concentration in CD of mM	Concentration of Compound I in $\mu\text{g/ml}$
10	120.6
25	526.7
50	1529.2
75	3012.4
100	4677.2
150	8317.6
200	11962.5

Table 4:
Solubility of Compound I in the presence of γ CD

Concentration in CD of mM	Concentration in Compound I of $\mu\text{g/ml}$
10	1.3
25	3.2
50	11.4
75	13.6
100	19.5
200	29.1

5 Compound I forms complexes with all the studied cyclodextrins because an increase in solubility is observed. It can also be directly observed that the complex formed between compound I and the Rameb increases the aqueous solubility of compound I considerably and completely unexpectedly. This observation is also true for the HP- β -CD. Table 5 summarizes the solubility results obtained for each cyclodextrin at the maximum concentration tested. The increase in solubility is
10 calculated compared to the solubility of compound I in water (in the absence of cyclodextrin) which has been determined at 0.56 $\mu\text{g/ml}$.

Based on these results, phase-solubility diagrams were constructed according to Higuchi, T., and Connors, K.A., Advances in Analytical Chemistry and Instrumentation 4 (1965) 117-212.

Table 5:

Maximum increase in solubility of Compound I obtained for each cyclodextrin

	Maximum concentration used in mM for each CD	Maximum solubility in $\mu\text{g/ml}$	Increase in solubility
βCD	4	6.8	14 x
HP βCD	200	4962	925 x
RM βCD	200	11926.5	23925 x
γCD	200	29.1	58 x

Table 6:

CD	Stoichiometry	$K_{1:1}[\text{M}^{-1}]$	$K_{1:2}[\text{M}^{-1}]$
βCD	1:1; 1:2	2092	-
γCD	1:1	346	-
HP- βCD	1:1; 1:2	12575	14.4
RM βCD	1:1; 1:2	27595	22.88

5

The high values of $K_{1:1}$ suggest that, in purified water, the cavity of the $\beta\text{-CD}$ derivatives accommodates very well the molecular portion of compound I involved in the inclusion. From 0 to 4 mM βCD concentration, the solubility of compound I increases and reaches a plateau up to 8 mM βCD . Concentrations of above 8 mM βCD form an additional complex of 1:2 stoichiometry (Cpd. I : βCD) with a lower solubility (1.5 $\mu\text{g/ml}$). The phase diagram obtained therefore is an A_L diagram. For $\gamma\text{-CD}$, HP- βCD and RM βCD , an A_p type diagram is obtained. The calculated stability constant of 346M^{-1} indicates that the cavity of the CD is too large to obtain sufficient interactions.

10

15

Compound I has different solubility when it is in the complex form or not. For example, compound I shows a good solubility in acetonitrile ($\pm 700\mu\text{g/ml}$) while the HP- $\beta\text{-CD}$ and the compound I-CD complex is insoluble in this solvent. In these conditions the included drug remains trapped and becomes insoluble in the solvent. This technique of differential solubility between compound I in the free or in the complexed form can be applied to evaluate the percentage of complexation.

20

Example 3

Solubility of various trioxypyrimidines in the presence of HP β CD

Solubility is investigated according to Examples 1 and 2. The results are shown in Table 7.

5

Table 7:

Concentration HP β CD [mM]	Concentration [mg/ml]			
	Compound III	Compound II	Compound IV	Compound V
10	0.8	1.3	0.1	0.8
25	2.4	3.4	0.2	2.4
50	3.1	3.8	0.9	5.4
100	6.1	5.9	2.7	6.6
200	9.5	9.3	7.9	9.9

Example 4

NMR studies

10

15

20

DIME- β -CD solutions were prepared in D₂O at 10 mM concentration. As water solubility of compound I is too low, spectra of compound I alone could not be performed in D₂O. For assignment of the protons, NMR spectra of compound I were performed in DMSO. All NMR experiments were performed on a Bruker DRX500 spectrometer operating at 500 MHz for proton. The temperature was set to 298K. Calibration was achieved using the residual resonance of the solvent as secondary reference of HDO. For T-ROESY experiments, a 300 msec mixing time was used. All processing were done on Silicon Graphics INDY data stations using WINNMR program for Bruker. The comparison between compound I NMR spectra alone and in presence of an excess of DIME β CD allows to notice that the signals corresponding to H-3 and H-5 protons are shifted up field. This shift constitutes a proof of the inclusion. The T-ROESY spectra analysis demonstrates the inclusion of compound I in the CD cavity. Two different parts of the molecule can fit into CDs cavity.

Example 5

Molecular modeling studies

Molecular modeling calculations have been performed with Gaussian 94 using the POBRON crystallographic structure of β -CD Of the Cambridge Data Base. Two extreme spatial conformations of compound I were calculated. The results obtained show the inclusion is energetically feasible and very stable. This stability can be explained by the formation of hydrogen bounds between the oxygen and the proton of the nitrogen of the barbituric nucleus and the alcohols situated at the outside of the CD. All the pharmaceutical compositions including compound I and a cyclodextrin (preferably β CD, γ CD and their synthetic derivatives) either in complexes form, or in association, are anticipated in the framework of the invention, whatever their form and their therapeutic application. In fact, even if compound I and the CDs are not in the form of complex in the formulation, this one is susceptible to be formed in situ.

Example 6

Pharmaceutical compositions

Different compositions of formulations are given for example non-exhaustively.

A preferred example for an injectable formulation is:

- HP- β CD 200 mM; Compound I 1 mg/ml; Sterile water for injection q.s.

For 25 ml of solution:

a) Preparation of the solution:

Weigh 6.77 g of HP β CD (4.2 % of H₂O) and dissolve them in 25 ml of water by injection. Add 25 mg of compound I and heat in a water bath until the latter is completely dissolved. Sterilize the solution by filtration.

b) Characteristics of the solution:

The solution osmolality is 308 mOs/kg. The pH is 7.2.

The concentration of compound I and/or of CD can be modified in function of the requirements. It is preferred to adjust the tonicity by addition of NaCl.

A preferred formulation for nebulization is:

For 200 ml of solution:

- | | | |
|---|--|----------------------|
| | - Compound I | 0.1 g (MW: 485) |
| 5 | - HP β CD exempt from pyrogenic | 20.15 g (MW: 1,300) |
| | - Sodium chloride | 1.42 g (isotonicity) |
| | - Pyrogen-free, sterile, purified water, | q.s. ad 200 ml |
- a) Weigh 20.15 g of HP β CD exempt from pyrogenic (3.2 % H₂O, ROQUETTE)
10 and dissolve them in 100 ml of purified water.
- b) Weigh 0.1 g of compound I, and 1.42 g of sodium chloride and add them to
solution (a) by energetically stirring so as to dissolve them.
- c) Complete with water so as to obtain 200 ml of solution.
- d) Sterilize by filtration through a 0.22 μ m polypropylene membrane.

15

Example 7

In vivo experiments (inhibition of angiogenesis)

20

In order to study the potential effects of the complex Compound I-cyclodextrine, a model of neovascularisation has been used. An aorta ring is cut and placed in a culture medium. This culture medium containing either:

- no active principle
- the complex compound I-cyclodextrins (final concentration 10⁻⁶M, 10⁻⁷M)
- compound I dissolved in DMSO with the help of DMSO (final concentration 10⁻⁶, 10⁻⁷M).

25

In the absence of the matrix metalloproteinase inhibitor compound I, the formation of new vessels (angiogenesis) is observed. In the presence of compound I alone, dissolved in DMSO, or in the form of inclusion complex in cyclodextrin, angiogenesis is inhibited significantly.

Example 8

Use of formulations containing COMPOUND I and HP β CD for therapy of allergen-induced airway inflammation and bronchial hyperresponsiveness in a mouse model of asthma

5 Materials

HP- β -CD (degree of substitution = 0.64) originates from Roquette (France). Apyrogenic phosphate buffered saline (PBS) was purchased from Bio-Wittaker (Verviers, Belgium) and methacholine from Sigma-Aldrich (Germany). All other materials were of analytical grade. Sterile water for injection was used throughout this study. Sterile, apyrogenic and isotonic CD solutions were prepared at 20, 50 and 75 mM. A commercially available fluticasone solution for inhalation (Flixotide® 1mg/ml) was purchased from Glaxo-Smithkline (Genval, Belgium)

Sensitization, allergen exposure and therapeutic protocols

In order to study the modulation of airway inflammation by intraperitoneal injection of Compound I, mice were sensitized with 10 μ g ovalbumin alumin-adsorbed (aluminject, perbio, Erembodegem, Belgium) injected intraperitoneally at days 0 and 7 and were subsequently exposed to ovalbumin (OVA) 1% or PBS aerosols for 30 minutes from day 21 to 24. Intraperitoneal injections were performed 30 min before OVA inhalations. The different injected formulations were: cremophor 10%-DMSO 10%-PBS 80%- Compound I 30 mg/kg (suspension); cremophor 10%-DMSO 10%-PBS 80%- Compound I 3.75 mg/kg (solution); HP β CD 200 mM Compound I 7.5 mg/kg (solution); HP β CD 200 mM. All results were compared to mice sensitized with OVA and exposed to PBS and OVA treated with PBS injected intraperitoneally. In order to study the modulation of airway inflammation by inhaled Compound I, mice were sensitized as described previously. Two protocols referred to as short exposure challenge and long-term exposure challenge were used. In the short exposure challenge, mice were exposed to aerosols of Compound I- complex at concentrations of 0.03 and 0.3 mg/ml of active compound in aqueous solution of from day 21 to 27 during 30 min in a Plexiglas exposure chamber (30 x 20 x 15 cm). Mice were exposed to OVA aerosols 30 minutes after the Compound I inhalation from day 23 to 27. In the so called long-term inhalation challenge, mice were exposed to aerosols of Compound I at concentrations of 0.03 and 0.3 mg/ml complexed with HP β CD in an aqueous

solution during 30 min five days odd weeks and to OVA aerosols 3 days odd weeks for 11 weeks. No inhalations were performed during even weeks.

5 The aerosol were produced by using an ultrasonic nebuliser SYSTAM (Système Assistance Medical, Le Ledat, France), the vibration frequency of which is 2.4 MHz with variable vibration intensity and ventilation levels. Vibration intensity was fixed in position 6 and the ventilation level was $25(\nu_{1/2})$ l/min.

Airway responsiveness measurement

10 Twenty-four hours after the last allergen exposure, the bronchial hyper responsiveness was determined by measuring the Penh using a barometric plethysmograph as proposed by Hamelmann, E., et al., Am. J Respir. Crit. Care Med. 156 (1997) 766-775). The Penh was measured at baseline and 5 min after the inhalation of increasing doses (25, 50, 75 and 100 mM) of methacholine (Mch).

Bronchoalveolar lavage (BAL) and histology

15 Immediately after the assessment of airway responsiveness, mice were sacrificed and 1 ml of PBS free of ionised calcium and magnesium but supplemented with 0.05 mM sodium EDTA was instilled 4 times via a tracheal cannula and recovered by gentle manual aspiration. The recovered bronchoalveolar lavage fluid (BAL) was centrifuged (1800 rpm for 10 min at 4°C). The cell pellet was washed twice and finally resuspended in 1 ml of PBS. A total cell count was performed in a Thoma
20 chamber and the differential cell counts on at least 400 cells were performed on cytocentrifuged preparations (Cytospin 2; Cytospin, Shandon td., Runcorn, Cheshire, U.K.) using standard morphologic criteria after staining with Diff-Quick (Dade, Germany). After BAL, the thorax was opened and the left main bronchus was clamped. The left lung was excised and frozen immediately in liquid N₂ for
25 protein chemistry and mRNA extraction while the right lung was processed for histology. As previously described (Cataldo, D.D., et al, Am. J. Pathol. 161 (2002) 491-498), the right lung was infused with 4% paraformaldehyde and embedded in paraffin. Sections of 5 µm thickness from all lobes were stained with haematoxylin and eosin. The extent of peribronchial infiltrates was estimated by an inflammation
30 score. Slides were coded and the peribronchial inflammation was graded in a blinded fashion using a reproducible scoring system described elsewhere (Cataldo, D.D., et al., Am. J. Pathol. 161 (2002) 491-498). A value from 0 to 3 per criteria was

adjudged to each tissue section scored. A value of 0 was adjudged when no inflammation was detectable, a value of 1 for occasional cuffing with inflammatory cells, a value of 2 when most bronchi were surrounded by a thin layer (1 to 5 cells) of inflammatory cells and a value of 3 when most bronchi were surrounded by a thick layer (>5 cells) of inflammatory cells. As 5-7 randomly selected tissue sections per mouse were scored, inflammation scores could be expressed as a mean value per animal and could be compared between groups. Another score referred to as tissue eosinophil infiltration score, specifically reflecting the amounts of eosinophils infiltrating the bronchial walls, was measured as follows: after a congo red staining, seven bronchi were studied per mouse. The eosinophils were counted around the bronchi within the limits of the airway wall, the perimeter of the epithelial basement membrane was measured and the results were expressed as number of eosinophils/mm of basement membrane. The left lung was snap frozen in liquid nitrogen and crushed using a Mikro-Dismembrator S (Braun Biotech International, Melsungen, Germany) and the extracts stored at -80°C before studied. Kidneys were excised and paraffin embedded, sections of $5\text{ }\mu\text{m}$ were stained by haematoxylin and eosin. Blood was sampled by cardiac puncture and serum was stored at -80°C until analysis were performed.

All in vivo manipulations were approved by the local Veterinarian Ethics Committee.

Intraperitoneal injection of Compound I

The intraperitoneal injection of Compound I (either solution or precipitate) lowered the allergen-induced airway eosinophilic inflammation in BAL at doses of 3.75 to 30 mg/kg when compared to placebo (Figure 4a). At the same doses, the peribronchial inflammation scores were also significantly lowered by Compound I with an equal efficacy of all tested formulations (Figure 4b). The tissue eosinophil infiltration score was significantly lowered by the intraperitoneal injection of Compound I at doses of 7.5 and 25 mg/kg.

Inhalational exposure to Compound I and Compound I- HP β CD complexes

The intrinsic activity of Compound I was firstly assessed as a topically active anti-inflammatory agent by using a solution of Compound I 40mg/ml in pure DMSO in a short-term exposure. When compared to the inhalation of DMSO alone, the

inhalation of this formulation led to a significant decrease of BAL eosinophils ($p < 0.005$), peribronchial inflammation scores ($p < 0.01$), as well as bronchial hyperresponsiveness ($p < 0.05$).

5 In the short-term exposure protocol, we assessed the effects of HP- β -CD
Compound I complexes containing formulations on the airway inflammation and
hyperresponsiveness. The effects of inhalation of Compound I- HP β CD complex
containing formulations were compared with those of placebo (PBS) or fluticasone
(1mg/ml) used as reference therapy. Inhalation of those formulations containing
10 Compound I at doses of 0.03 and 0.3 mg/ml induced a significant decrease in
eosinophilic inflammation in BAL in an extent comparable to that of fluticasone
when compared to placebo ($p < 0.0001$) (Figure 5a). Peribronchial inflammation
scores were also lowered when compared to placebo ($p < 0.0001$) (Figure 5b), as well
as the tissue eosinophil infiltration score ($p < 0.01$) (Figure 5c).


15 After long term allergen exposure, BAL eosinophilia was significantly decreased
after treatment by inhalation of Compound I- HP β CD containing formulations
($p < 0.001$) in the same extent as that of fluticasone (Figure 6a). The peribronchial
inflammation score was also significantly decreased by inhalation of Compound I-
HP β CD containing formulations as well as by fluticasone ($p < 0.0001$) (Figure 6b).
The tissue eosinophil infiltration score was also decreased after treatment by
20 Compound I inhalation in an extent comparable to the fluticasone treated mice
($p < 0.01$) (Figure 6c).

List of References

- Bergers, G., et al., Nat. Cell Biol. 2 (2000) 737-744
Carmeliet, P., et al., Nat. Genet. 17 (1997) 439-444
Carstanjen, D., et al., Transfusion 42 (2002) 588-596
5 Cataldo, D.D., et al., Am. J. Pathol. 161 (2002) 491-498
Chang, C., and Werb, D., Trends Cell Biol. 11 (2001) S37-43
Dong, Z., et al., Cell 88 (1997) 801-810
Egeblad, M., and Werb, Z., Nat. Rev. Cancer 2 (2002) 161-174
EP 0 869 947
10 Fabbri, L.M., and Hurd, S.S., Eur. Respir. J. 22 (2003) 1-2
GINA Workshop Report, Global Strategy for Asthma Management and
Prevention (NIH Publication No. 02-3659)
Grams, F., et al., Biol. Chem. 382 (2001) 1277-1285
Hamelmann, E., et al., Am. J. Respir. Crit. Care Med. 156 (1997) 766-775
15 Higuchi, T., and Connors, K.A., Advances in Analytical Chemistry and
Instrumentation 4 (1965) 117-212
Holmbeck, K., et al., Cell 99 (1999) 81-92
Lund, L.R., et al., EMBO J. 18 (1999) 4645-4656
Manes, S., et al., J. Biol. Chem. 274 (1999) 6935-6945
20 Overall, C.M., and Lopez-Otin, C., Nat. Rev. Cancer 2 (2002) 657-672
Remington's Pharmaceutical Sciences, 16th ed., 1980, Mack Publishing Co.,
edited by Oslo et al.
Shapiro, S.D., Curr. Opin. Cell Biol. 10 (1998) 602-608
US 6,110,924
25 US 6,242,455
Vu, T.H., et al., Cell 93 (1998) 411-422
WO 01/25217
WO 97/23465
WO 98/58915

Patent Claims

1. A trioxypyrimidine-cyclodextrin complex formed of a trioxypyrimidine derivative or a salt thereof and a water-soluble cyclodextrin, wherein the trioxypyrimidine derivative is represented by formula (I):



wherein

R^1 is C_3 - C_{20} alkyl, which may optionally be interrupted once or several times by -S-, -O- or -NH-; or

a group W-V, wherein

W is a chemical bond or phenyl; and

V is phenyl, phenyloxy, phenylthio, phenylsulfinyl, phenylsulfonyl or phenylamino, which phenyl moieties may be unsubstituted or substituted once or several times by halogen, hydroxy, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_1 - C_6 -alkylthio, C_1 - C_6 alkylsulfinyl, C_1 - C_6 -alkylsulfonyl, cyano, nitro or C_1 - C_6 -alkylsulfonyl; and

R^2 is C_1 - C_{10} alkyl, which alkyl group is unsubstituted or substituted one or two times by hydroxy or amino and may optionally be interrupted once or several times by -S-, -O- or -NH-;

a benzoyl group, which may be unsubstituted or substituted once or several times by halogen, hydroxy, nitro, C_1 - C_6 -alkoxy, C_1 - C_6 -alkylamino, C_1 - C_6 -alkylthio, C_1 - C_6 -alkylsulfinyl, C_1 - C_6 -alkylsulfonyl, amidosulfonyl, C_1 - C_6 -alkylamid sulfonyl, bis- C_1 - C_6 -alkylamid sulfonyl;

a heteroaromatic acyl group; or

a phenyl- or heteroaryl group, which are unsubstituted or substituted once or several times by halogen, hydroxy, C₁-C₆-alkoxy, C₁-C₆-alkylamino, C₁-C₆-dialkylamino, cyano, C₁-C₆-alkyl, C₂-C₆ alkenyl, C₂-C₆-alkinyl, C₁-C₆-acyl, C₁-C₆-alkylthio, C₁-C₆-alkylsulfonyl, C₁-C₆-alkylsulfinyl, C₁-C₆-alkylaminocarbonyl, aminocarbonyl, C₁-C₆-alkylamidossulfonyl, amidosulfonyl, bis-C₁-C₆-alkylamidossulfonyl, nitro, C₁-C₆-alkoxycarbonyl, carboxy.

2. A trioxypyrimidine-cyclodextrin complex according to claim 1, wherein the trioxypyrimidine derivative is

5-Biphenyl-4-yl-5-[4-(4-nitro-phenyl)-piperazin-1-yl]pyrimidine-2,4,6-trione;

5-(4-Phenoxy-phenyl)-5-(4-pyrimidin-2-yl-piperazin-1-yl)-pyrimidine-2,4,6-trione;

5-[4-(4-Chloro-phenoxy)-phenyl]-5-(4-pyrimidin-2-yl-piperazin-1-yl)-pyrimidine-2,4,6-trione ;

5-[4-(3,4-Dichloro-phenoxy)-phenyl]-5-(4-pyrimidin-2-yl-piperazin-1-yl)-pyrimidine-2,4,6-trione;

5-[4-(4-Bromo-phenoxy)-phenyl]-5-(4-pyrimidin-2-yl-piperazin-1-yl)-pyrimidine-2,4,6-trione

or a salt thereof.

3. A trioxypyrimidine-cyclodextrin complex according to any one of claims 1 to 2, wherein the water-soluble cyclodextrin is β -cyclodextrin.
4. A trioxypyrimidine-cyclodextrin complex according to any one of claims 1 to 2, wherein the water-soluble cyclodextrin is hydroxypropylated cyclodextrin.

5. A trioxypyrimidine-cyclodextrin complex according to any one of claims 1 to 2, wherein the water-soluble cyclodextrin is random methylated cyclodextrin.
6. A trioxypyrimidine-cyclodextrin complex according to any one of claims 1 to 2, wherein the water-soluble cyclodextrin is sulfobutyl- β -cyclodextrin.
- 5 7. A trioxypyrimidine-cyclodextrin complex according to any one of claims 1 to 2, wherein the water-soluble cyclodextrin is γ -cyclodextrin.
8. A pharmaceutical formulation containing a trioxypyrimidine-cyclodextrin complex as defined in any one of claims 1 to 7.
9. A pharmaceutical formulation according to claim 8 containing a
10 pharmaceutically acceptable additive.
10. A process for preparing a complex selected from the group consisting of α -cyclodextrin and trioxypyrimidine, β -cyclodextrin and trioxypyrimidine, and γ -cyclodextrin and trioxypyrimidine comprising mixing said cyclodextrin and trioxypyrimidine in a molecular ratio between 2:1 and 540:1 for at least
15 seven days at a temperature between 20 and 80°.

0 1. April 2004

Abstract

A trioxypyrimidine-cyclodextrin complex formed of a trioxypyrimidine derivative or a salt thereof and a water-soluble cyclodextrin derivative has improved solubility.

Fig. 4a




Fig. 4b




Fig. 3





Fig. 2



01. April 2004


Fig. 1

Fig. 5c




Fig. 5a




Fig. 5b




Fig. 6a




Fig. 6b




Fig. 6c

